# On the Clear Evidence of the Risks to Children from Smartphone and WiFi Radio Frequency Radiation

Professor Tom Butler | University College Cork | tbutler@ucc.ie

Children's health and well-being is under significant threat from everyday digital technologies, as the past 15 years have seen the proliferation of microwave Radio Frequency Radiation (RFR) devices in the home, school and society. The safety standards for such devices—smartphones, tablets etc.—were based on the proven thermal or heating effects of microwave devices in adults, not children. Scientists in the life sciences have long been aware of equally harmful non-thermal effects. However, physicists and engineers have operated on the theory that nonionizing RFR could not directly damage human DNA and lead to cancer, as it was far less powerful than ionizing radiation (xrays, nuclear etc.). That theory has been solidly and roundly refuted, as this paper illustrates. Cancer risks aside, research studies have demonstrated that low-intensity RFR elicits a range of pathophysiological conditions in experimental animals and humans. This is why parents, educators and governments should be alarmed and take immediate and appropriate action. In an Irish, and indeed international context, there is clear ignorance on such matters. In response, this paper aims to inform by presenting clear evidence, in a balanced manner, of the risks to children based on proven scientific theories and empirical evidence. The paper concludes by offering practical advice on how the risks to children, and indeed adults, can be minimised.





Professor Tom Butler University College Cork tbutler@ucc.ie

> "The level of proof required to justify action for health protection should be less than that required to constitute causality as a scientific principle"

> > Professor Rainer Frentzel-Beyme MD

#### **Contents**

Abstract3
Introduction3
Proof of the Potential Toxicity and Carcinogenicity of RFR4
What are the Biological Mechanisms that Produce Ill-health in Children and Adults?7
Evidence that Microwave RFR Promotes the Development of Existing Cancers9
Why are Existing Standards Unsafe?9
What do Insurance Companies and Regulators and Telecoms Operators have to say about the Risks?11
Discussion12
Conclusions14
About the Author17
References17

#### Abstract

Children's health and well-being is under significant threat from everyday digital technologies, as the past 15 years have seen the proliferation of microwave Radio Frequency Radiation (RFR) devices in the home, school and society. The safety standards for such devices smartphones, tablets etc.-were based on the proven thermal effects microwave devices in adults, not children. Scientists in the life sciences have long been aware of equally harmful non-thermal effects. However, physicists and engineers have operated on the theory that non-ionizing RFR could not directly damage human DNA and lead to cancer, as it was far less powerful than ionizing radiation (x-rays, nuclear etc.). That theory has been solidly and roundly refuted, as this paper illustrates. Cancer risks aside, research studies have demonstrated that low-intensity RFR elicits a range of pathophysiological conditions in experimental animals and humans. This is why parents, educators and governments should be alarmed and take immediate and appropriate action. In an Irish, and indeed international context, there is clear ignorance on such matters. In response, this paper aims to inform by presenting clear evidence, in a balanced manner, of the risks to children based on proven scientific theories and empirical evidence. The paper concludes by offering practical advice on how the risks to children, and indeed adults, can be minimised.

#### Introduction

Exposure of humans to non-ionizing radio frequency radiation (RFR) has increased dramatically over the past 20 years, particularly where children are concerned. Epidemiological and experimental research reports increased risk of pathophysiological conditions with current exposures to Smartphone and WiFi RFR that include: cellular DNA damage, leading to a range of cancers; sperm and testicular damage leading to

male infertility; neuropsychiatric conditions, including post-natal neural development, and learning and cognitive problems; and melatonin reduction leading to sleep disruption; among others. There is a considerable body of evidence on the harmful effects of electromagnetic pollution that should have the World Health Organization and governing (WHO) establishing guidelines to protect public health, as they have with environmental pollution. In light of recent scientific evidence, in May 2015 over 200 eminent scientists have launched an international appeal to the United Nations and the WHO based on the conviction that there is a real and present danger to children, in particular, by what they consider outdated industry standards in relation to microwave radiation.

On November 1st 2018, the final report of a 10-year \$30m comprehensive study US National Institute by Environmental Health Sciences' National Toxicology Program (NTP) confirmed that radio frequency radiation (RFR) from Smartphones caused cancer in animals.<sup>1</sup> That study clearly refutes the long-held theory that non-ionizing radiation, such as RFR, cannot cause cancers or lead to other effects on the health and wellbeing of humans. The findings of this study opens an ethical Pandora's Box for mobile phone companies and BigTechs such as Apple, Facebook, Google and others, as the use of microwave RFR technologies underpin their business models. Such technologies are now ubiquitous in society, whether it is in the home, classroom, workplace, or in transport systems. The fact that they might pose a real risk to the health and well-being of users and particularly children was never considered. This creates a dilemma for parents and educators, as the evidence on the risks to human health and well-being associated with widespread indiscriminate exposure to RFR is clear and unambiguous, with children being particularly at risk.

This paper begins by reporting on a ground-breaking study by the National Toxicology Program's (NTP) at the U.S. Department of Health and Services. The final report was issued on November 1st 2018 and in the press Dr. John Bucher, Senior release Scientist, at the National Toxicology Program stated, "We have concluded that there was clear evidence that male rats developed cancerous heart tumors called malignant schwannomas. occurrence of malignant schwannomas in the hearts of male rats is the strongest study." <sup>2</sup> cancer finding in our Categorising the major findings as "clear evidence" is significant as this is the highest burden of proof in a scientific study by the NTP. It employs 4 levels of evidence. Other findings categorised as Some Evidence (brain tumours such as glioma and adrenal gland tumours) and Equivocal (cancers of the prostate and pituitary glands). None of the findings were at level 4, No Evidence. The paper discusses these findings in the context of previous research. It then presents the biological and cellular mechanisms found to be responsible for these effects. The paper presents evidence that then promotes the development of existing cancers and examines why the existing safety standards are not fit for purpose. Next explored are the perspectives of insurance companies and telecommunications operators on the risks.

### Proof of the Potential Toxicity and Carcinogenicity of RFR

In 1999, the US Food and Drua Administration's (FDA) Center for Devices and Radiological Health commissioned the National Toxicology Program study on the potential toxicity and carcinogenicity of RFR.<sup>3</sup> The FDA's concerns followed the emergence and widespread use of first generation cell phone devices in the early 1980s and second generation (2G) systems in the 1990s. The health focus and associated

safety standards were, and still are, centred on the thermal effects (i.e. heating of tissues from microwaves) and not on the non-thermal effects. To be sure, the findings of extant research at the time were mixed, with no clear evidence either way of potential negative health implications of low-intensity RFR, especially where cancer was concerned.4 In 2011 the IARC classified WiFi and microwave radiation from cordless and mobile phones as a *possible* Class 2B carcinogen. While the findinas epidemiological studies have been debated, and chiefly focus on the longterm development of brain tumours, a recent review of such studies is unequivocal and states that "[m]obile phone radiation causes brain tumors and should be classified as a probable human carcinogen (2A)" by the WHO's International Agency for Research on Cancer (IARC).5

The NTP study is not the first of its kind—it confirms the findings of previous research on the links between RFR exposure and various cancers—it is the most comprehensive, however. Since 1999 when the FDA flagged the issue of non-thermal potential effects microwave radiation, a wealth of new epidemiological experimental and research demonstrated the very real biological effects of RFR on the brain, nervous systems, hearts and testes of mammals, including humans. Cancers aside, many of these studies consistently report a range of side-effects in humans, from sleep deprivation and headaches, to neurological damage, and learning disorders. Consequently, in 2014, the California Medical Association (CMA) 6 stated that "peer reviewed research has demonstrated adverse biological effects of wireless EMF [i.e. RFR] including single and double stranded DNA breaks, creation of reactive oxygen species, immune dysfunction, cognitive processing effects, protein stress synthesis in the brain, altered brain development, sleep and memory disturbances, ADHD, abnormal behavior,

sperm dysfunction, and brain tumors." The CMA were following the lead of the American Academy of Paediatrics, which in 2013 petitioned the US Federal Communications Commission (FCC) and to the Food and Drug Administration (FDA) to reassess safety standards to RFR in order to "protect children's health and well-being throughout their lifetimes and reflect current use patterns."

The final peer-reviewed findings of the NTP study were released on November 1<sup>st</sup> 2018. Given the significance of the findings there was a muted response from the press. Coverage from the New York Times argued that the focus on 2G and 3G technologies somehow weakened the study's findings.8 This is a spurious argument, as 4G Smartphones are backward compatible with 2G and 3G. worryingly the International Commission on Non-Ionizing Radiation Protection (ICNIRP) decided that the findings did not provide a reason to revise current (i.e. 20-year-old) RFR standards. However, exposure Ronald Melnick rebutted the ICNIRP analysis stating it contained several false and misleading statements. 9 Dr Fiorella Belpoggi, Director of the Cesare Maltoni Cancer Research Center of the Ramazzini Institute, which had recently conducted separate research that reproduced the findings of the NTP Study, also took issue ICNIRP—"We with the scientists, our role is to produce solid evidence for hazard and risk assessment. Underestimating the evidence from carcinogen bioassays and delays in regulation have already proven many times to have severe consequences, as in the case of asbestos, smoking and *vinyl chloride."* <sup>10</sup> In their study, Dr Belpoggi and her colleagues found an "increase in the incidence of tumors of the brain and heart in RFR-exposed Sprague-Dawley rats. These tumors are of the same histotype of those observed in some epidemiological studies on cell phone users." 11 The NTP study also reported that DNA damage (strand breaks) was significantly increased in the

brains of rats and mice exposed to RFR. The findings also reported reduced birth weights of rat pups whose mothers were exposed to RFR, in addition to cardiomyopathy of the right ventricle in the rats studied.<sup>12</sup>

After more than 20 years of widespread cell phone use, one would expect to see a rise in cancers, particularly brain tumours. A research review of the incidence of glioblastoma multiforme tumours in England during 1995-2015 reported a "a sustained and highly statistically significant ASR [(incidence rate)] rise in glioblastoma multiforme (GBM) across all ages. The ASR for GBM more than doubled from 2.4 to 5.0, with annual case numbers rising from 983 to 2531. Overall, this rise is mostly hidden in the overall data by a reduced incidence of lower-grade tumours."

13 The study did not focus on RFR as the cause, so the findings must be considered 'open to interpretation' in this regard, as other environmental mechanisms cannot be ruled out. Another recent review study on epidemiological studies on brain and salivary gland tumours and mobile phone use found the cumulative evidence to be inconclusive, but indicated that cancers may have longer latency (i.e. greater than 15 years) and childhood use to be of concern. 14 In contrast, the French CERENAT study stated that, "Consistent with previous studies, we found an increased risk in the heaviest users, especially for gliomas." 15 While these studies emanate from natural scientists, two social scientists found that "that mobile phone subscription rates are positively and statistically significantly associated with death rates from brain years later. 15-20 As cancer falsification test, we find few positive associations between mobile phone subscription rates and deaths from rectal, pancreatic, stomach, breast or lung cancer or ischemic heart disease."16

Commenting on the general issue of a rise in brain tumours, cancer epidemiologist Geoffrey Kabat, argues that the risk of brain cancer in the US is

low, 6 in every 100,000, compared to 100,000 for breast 150 in every cancer. 17 He therefore implies that the cancer risk from RFR is overhyped, as epidemiological data can be open to various interpretations. Whichever way one looks at it, the incidence of such tumours is indeed low. But if recent studies are accurate, then this rate may more than double over a 20 year period<sup>11</sup> or increase in line with mobile phone use, 14 as RFR is now accepted as either a causal or a contributory mechanism in the occurrence of brain tumours and other cancers. However, what none of these studies take into account is that the number of RFR sources has increased dramatically throughout the home, school and work environments over the past 10 years, with WiFi routers, 2-4G enabled tablets, the proliferation of WiFi enabled devices and wearables, and the number of mobile phones per person.

There is also evidence that RFR from cell phones may be triggering breast cancer in young women who carry their devices on or near their breasts. 18 To compound matters even further, one of the significant findings of the NTP study was the presence of RFR promoted the growth of tumours caused by other carcinogens. 1,8,10,11 The findings of the cumulative bodv of research objective, and particularly disturbing where children are concerned. Rigorous experimental studies on laboratory rats have found that daily exposures to lowlevels of microwave radiation, such as that emitted by WiFi devices, causes significant biological changes in a range of major organs such as the brains, hearts, reproductive systems, and eyes of the rats being studied. 19

All this has profound implications for the increasing numbers of children and adolescents exposed to RFR on a daily basis. And the risks to children are considerable: "Because cells are rapidly dividing and organ systems are developing during childhood and adolescence, exposure to carcinogens

during these early life stages is a major risk factor for cancer later in life. Because young people have many expected years of life, the clinical manifestations of cancers caused by carcinogens have more time in which to develop during characteristically long latency periods." 20 A recent study demonstrated that in a child's brain the hippocampus and hypothalamus absorb 1.6-3.1 times the microwave energy of an adult brain. The absorption rate is 2.5 times higher than an adult's where a child's cerebellum is concerned. The same study found that the bone marrow in a child's skull absorbs microwave radiation at a level 10 times greater than that of an adult.<sup>21</sup> In addition, a child's eyes absorb higher levels of microwave radiation than adults. 22 If, as the latest scientific evidence indicates, low-level microwave radiation poses a health risk, and if safety standards are outdated, then it is logical to assume that children are at significant risk from any device radiating microwave radiation.<sup>23</sup>

Dr Christopher J. Portier, Associate Director, National Institute Environmental Health Sciences and Director, Office of Risk Assessment Research, co-authored an article with Dr Wendy Leonard in Scientific American, following the initial release of the NTP study findings in 2016. They conclude that, "Cellphones probably cause cancer if the exposure is close enough, long enough, and in sufficient magnitude. We don't yet know the risk for a given level of exposure in humans. We need more data in this area, not only for cellphones, but for bluetooth devices, wifi and all the other RF-EMF devices out there. Until then, reduce your exposure whenever possible."24

Clearly, this is a complex matter, made even more so by the fact that there was no hope of a paradigm change, until the 'smoking gun' provided by the NTP study removed any doubt that RFR can act directly, via identified mechanisms, to induce tumours in biological organisms




Figure 1 Mechanisms and Pathways to Pathophysiological Effects (Reproduced from Pall 2018)

exposed to radiation levels within those permitted by existing standards and to which users are typically exposed. This should stimulate a reassessment of the risks in relation to all RFR use, particularly children. "The level of proof required to justify action for health protection should be less than that required to constitute causality as a scientific principle".<sup>25</sup> We are far beyond that level of proof where RFR is concerned.

### What are the Biological Mechanisms that Produce Ill-health in Children and Adults?

While the direct effects of certain carcinogens are widely acknowledged, research illustrates that "carcinogens may also partly exert their effect by generating reactive oxygen species (ROS) during their metabolism. Oxidative damage to cellular DNA can lead to mutations and may, therefore, play an important role in the initiation progression of multistage carcinogenesis...Elevated levels of ROS and down regulation of ROS scavengers and antioxidant enzymes are associated with various human diseases including various cancers. ROS are also implicated diabetes and neurodegenerative diseases"26. Research on mobile phone RFR and WiFi pulsed microwave signals has also demonstrated that they produce elevated levels of reactive oxygen species (ROS) which in turn cause oxidative stress in cell. 27,28, 29 Oxidative stress is caused by an imbalance between ROS and the counter effects of antioxidants that help detoxify and repair biological systems. Thus, the body normally employs antioxidant defence mechanisms to counter ROS and help avoid diseases such as cancer, which are trigged by oxidative stress and its tendency to cause strand breaks in a cell's DNA.

A raft of studies indicate that a chain of biological mechanisms produces the observed negative health outcomes in laboratory animals and humans. Martin Pall, Professor Emeritus of Biochemistry and Basic Medical Sciences, Washington State University points to the role of voltage-gated calcium channel (VGCC)

activation, which is triggered by RFR sources such as 2-4G and Wifi, as being one of primary causal mechanisms.<sup>30</sup> In his review published in 2018, he cites over 120 empirical research papers in support of his thesis. There is, therefore, a cumulative body of evidence which refutes the proposition that RFR has no biological effects, other than thermal effects on tissue. Professor Pall's earlier 2013 review paper cites 22 research studies that specifically point to the role played by VGCC activation. 31 The number of studies replicating experiments that corroborate this theory has grown significantly, while none appear to refute it. The figure 1 illustrates the posited mechanisms, pathways and outcomes. A detailed discussion is beyond the scope of this paper, however, several important mediating mechanisms and pathophysiological outcomes are now discussed.

A recent review of scientific studies concluded that relatively brief, regular, and also long-term use of microwave devices resulted in negative impacts on biological systems, especially the brain.<sup>32</sup> This review by Kesari et al. squarely highlights the role played by reactive oxygen species (ROS) as a key mechanism (generated by exposure to producing microwaves) in serious negative effects in living organisms. Exposure to ionizing radiation has been long known to disturb the balance between ROS and the antioxidants that neutralise them. Usually this imbalance results in a high probability that the subject will develop cancers and other chronic conditions. A wealth of studies illustrate, however, that nonnow ionizing radiation emitted from smart phones, cordless phones, WiFi, Bluetooth and other wireless technologies, such as those powering the Internet of Things (IoT) can severely disturb this balance also, by amplifying ROS, suppressing antioxidants, and increasing oxidative stress. There is substantial evidence that oxidative damage to cellular proteins,

lipids and DNA is at the root cause of many of the ill-effects of microwave RFR. Most worrying in all of this is that scientists have found that the mutagenic effects on the DNA of living cells occurs under low-levels of exposure to the pulsed microwave radiation found in most of these devices. The consequences for children are obvious, given their greater exposure levels and susceptibility to health ill-effects and also that their bodies are constantly growing and developing.<sup>33, 34</sup>

A recent study illustrates relatively low level of exposure required to produce adverse biological effects. Chauhan et al.8 published the results of their experiment on Wistar rats in 2016. The rats in this experiment were exposed to RFR at 25% of the normal level at the human ear and 15% of the level when carried for 2 hours per day for 35 days. Autopsies of the rats exposed to RFR revealed significantly high levels of ROS in the livers, brains and spleens of the exposed animals. In addition, histological changes were also found in brains, livers, testes, kidneys and spleens. In line with a wealth of other similar studies, the researchers concluded that the "results indicate possible implications of such exposure on human health." Earlier studies found that rat brains exposed to RFR exhibited an increase in sinale strand DNA breaks and chromosomal damage in brain cells. Thus it is beyond doubt that the substantial increase in ROS in living cells under RFR at low signal strength could be causing a broad spectrum of health disorders and diseases, including cancer, in humans and particularly in children. Certainly, recent studies have provided empirical evidence to support this theory.

Another recently discovered mechanism found to affect the arowth of multiforme in alioblastoma tumours protein. humans is the p53 Glioblastoma are the most common and most malignant of the glial tumours found in the brain and central nervous

system. These researchers studied 63 patients with this type of tumour and found that patients that used "mobile phones for ≥3 hours a day show a consistent pattern of increased risk for the mutant type of p53 gene expression the peripheral zone of glioblastoma, and that this increase was significantly correlated with shorter overall survival time." This is a significant finding. More worrying is a recent study conducted on the Swedish National Inpatient Register: "The main finding in this study was increasing rate of brain tumor of unknown type in the central nervous system." 36 The research being conducted by the 'Hardell Group' in Sweden, which is responsible for this study, has consistently demonstrated a link between mobile phone use and cancer. Two recent studies from the group confirm the link between RFR and cancers in humans. In the first, both mobile and cordless phones were associated with an increased the risk of glioma, a type of brain tumour. 37 It found that the "First use of mobile or cordless phone before the age of 20 gave higher OR [odds ratio] for glioma than in later age groups." Which indicates that children or teenagers are at significant risk. In the second, researchers found that the rise in thyroid cancers in Sweden was linked with increase in exposure to RF-EMF.<sup>38</sup> To be sure, epidemiological studies such as the latter are akin to looking for a needle in a haystack, and are criticised by some as being flawed, however their findings need to be viewed in a new light given the scientific evidence emerging from laboratory experiments such as the NTP study.

## Evidence that Microwave RFR Promotes the Development of Existing Cancers

One important recent finding is that RFR has cocarcinogenic effects. In research published in 2010, carcinogen-treated mice exposed to RFR demonstrated

significant tumour-promoting effects.<sup>39</sup> A study by Lerchl et al. in 2015<sup>40</sup> replicated the earlier study using higher numbers of animals in both the control and Lerchl et al. experimental groups. confirmed and extended the previous findings. They report that numbers of tumours of the lungs and livers in exposed animals in were significantly higher than in the control groups. They also reported significantly elevated lymphomas through RFR exposure. The hypothesized cocarcinogenic effects may have been "caused by metabolic changes due to exposure." Ιt is significant, extremely worrying, that tumourpromoting effects were produced "at low to moderate exposure levels (0.04 and W/kg SAR), thus well below exposure limits for the users of mobile phones." The authors conclude that their "findings may help to understand the repeatedly reported increased incidences of brain tumors in heavy users of mobile phones." The mechanisms presented in the previous section help explain why and how RFR exposures induce the observed findings in these and other studies.

#### Why are Existing Standards Unsafe?

The existing standards for mobile (2,3, & 4G) and WiFi are considered unsafe. The **US Federal Communications Commission** (FCC) mandates that "The safe limit for a mobile phone user is an SAR of 1.6 watts per kg (1.6 W/kg), averaged over one gram of tissue, and compliance with this limit must be demonstrated before FCC approval is granted for marketing of a phone in the United States." SAR is the Specific Absorption Rate. Expressed in Watts (a unit of electrical power) per kilogram of human tissue, SAR measures of the rate at which RFR energy is absorbed by the human body. In the testing procedures the FCC uses to certify that cell phones don't exceed the 1.6 W/kg SAR limit, the devices are tested 0.59 inches and 0.98 inches (1.5cm to 2.5cm) from the body. Hence,

smartphone manufacturers provide these quidelines buried in their safety information. If users operate their devices within these limits, which most do, they are in breach of the safe operating limits and are more at risk from both thermal and non-thermal effects. To make matters worse, 75% of smartphones regularly exceed safety limits as a recent correspondence between Washington DC law firm, Swankin & Turner, who sent a letter to the FCC indicates. The letter questioned whether the agency adequately enforced its cell phone radiation exposure limits.<sup>41</sup>

If a smartphone is on, but not being used for calls, text, or to browse online, it still communicates with cell phone base towers to maintain internet access. This allows app notifications, instant message texts, updates, and so on. So your phone is never off. Hence, when you carry it in your pockets or on a belt wallet, it's not being operated within the safe distance and the phone manufacturer is not liable. Note that the safety limits for cell phones and Wifi focus on thermal effects only. Remember also that non-thermal effects have been observed at much lower SAR levels from individual devices and also cellular base stations and WiFi router. 42 Russian scientist Dr. Yuri Grigoriev, Chairman of the Russian National Committee on Nonionizing Radiation Protection (RNCNIRP) points out that "National and international regulatory limits for radiofrequency radiation (RFR) exposure from cell phones and cell towers are outdated." 43 He argues that Western standards are inadequate to protect human health, in contrast with those in Russia, especially where the health of children is concerned. In Belpomme et al. study<sup>42</sup>, whose authors include cancer researchers, it is argued that "In spite of a large body of evidence for human health hazards from non-ionizing EMFs intensities that do not cause measureable tissue heating, summarized in an encyclopaedic fashion in the Bioinitiative Report

(www.bioinitiative.org), the World Health Organization (WHO) and governmental agencies in many countries have not taken steps to warn of the health hazards resulting from exposures to EMFs at low, non-thermal intensities, nor have they set exposure standards that are adequately health protective."

The industry safety standard for WiFi was established in 1996 by the FCC. It adopted the IEEE standard for safety levels with respect to human exposure to radio frequency electromagnetic fields, 3 kHz to 300 GHz, of 1991, which was based on that issued by the National Council on Radiation Protection (NRCP) in 1986. This standard covers only the thermal hazards from Radio Frequency Radiation (RFR). RFR is also known as non-ionizing microwave radiation. There long-standing belief physicists and electronics engineers that, unlike ionizing radiation such as X-rays, non-ionizing radiation such as RFR is not powerful enough to cause damage to human DNA. Non-ionizing microwave radiation could not, therefore, according to that theory be a cause of cancer or other non-thermal biological effects in humans. This theory has, however, been falsified, as indicated above.

hazards covered by the FCC standard are based on the specific absorption dose-rate (SAR) produces thermal effects in body tissue. As indicated SAR is typically measured in Watts/Kilogram. So, put simply, SAR estimates the amount of energy absorbed by a human body or part thereof when exposed to an RFR signal. While accurate, it chiefly focuses on thermal effects of RF-EMF. The FCC guidelines are based on a 4 W/Kg adverse level effect observed in laboratory animals. This excerpt from the Code of Federal Regulations (CFR 47/2.1093) is instructive:

"The SAR limits for general population/uncontrolled exposure are 0.08 W/kg, as averaged over the whole body, and a peak spatial-average SAR of

1.6 W/kg, averaged over any 1 gram of tissue (defined as a tissue volume in the shape of a cube). Exceptions are the parts of the human body treated as extremities, such as hands, wrists, feet, ankles, and pinnae, where the peak spatial-average SAR limit is 4 W/kg, averaged over any 10 grams of tissue (defined as a tissue volume in the shape of a cube). Exposure may be averaged over a time period not to exceed 30 minutes to determine compliance with general population/uncontrolled SAR limits."

Based on existing theories and research data, the FCC recognised the safety problems with WiFi and recommended that such devices are not operated less than 20 cm from the human body for 30 minutes. However, as far back as 2002, the Environmental Protection Agency (EPA) stated that the "FCC's exposure quideline is considered protective of from effects arising а thermal mechanism but not from all possible mechanisms. Therefore. generalisation by many the guidelines protect human beings from harm by any or all mechanisms is not justified"44. The EPA's reservations were justified, given research findinas published over the past 15 years that refute the theory that hazards were confined to thermal effects.



When is laptop not a laptop? When it's WiFienabled. Such devices must be 20 cm or 8" from the body— an adult body, that is.

We might add that at the time, the FCC never envisaged adults carrying WiFi enabled devices on their person, and

certainly never envisaged children using these devices on a regular basis. Note too that adults and children carry WiFienabled smartphones on their person, less than 1 cm from their bodies and well within the 20 (8") cm limit of safe operation. This is also true of WiFienabled WhatsApp and Skype calls. However, today such devices are in widespread daily use by children across developed countries. Furthermore, given the observable patterns of use, the 30minute maximum exposure is being breached on a regular basis by both adults and children. Given the scientific evidence, it is troubling to think that children are carrying or operating WiFi devices on or near their person, breaching the safety guidelines set by the FCC, and for periods much, much longer than 30 minutes.

Opposing views come from the BigTech and telecommunications companies, who like the tobacco lobby before them, are arguing that there is no danger in using WiFi technology or mobile phones. This view is based on the aforementioned belief that non-ionizing radiation such as microwaves are not powerful enough to cause damage to human DNA. However, Professor Martin Pall concludes "Repeated Wi-Fi studies show that Wi-Fi causes oxidative stress, sperm/testicular neuropsychiatric damage. effects changes, including EEG apoptosis, cellular DNA damage, endocrine changes, and calcium overload."30

### What do Insurance Companies and Regulators and Telecoms Operators have to say about the Risks?

In 2010 Lloyds of London<sup>45</sup> published a paper on the emerging risks of RFR. At the time it likened links between nonionizing radiation and cancer to that which exists between asbestos and cancer, indicating that time and more research would establish a causal link. In 2015, rumours spread across the web that Lloyds of London had stopped covering health risks associated with RFR

devices. However, it appears that the exclusion of RFR from insurance policies was issued by an individual underwriter, CFC Underwriting Ltd to the effect that: "The Electromagnetic Fields Exclusion (Exclusion 32) is a General Insurance Exclusion and is applied across the market as standard. The purpose of the exclusion is to exclude cover for illnesses caused by continuous long-term nonionising radiation exposure i.e. through mobile phone usage." It was reported that this exclusion applied to insurance cover for architects and engineers in Canada, following health concerns centering on a programme to install Wi-Fi in all British Columbian schools without parents' consent. Lloyd's 2010 report predated the IARC's decision in 2011 to classify RFR as a Class 2B carcinogen. As research on the health risks of RFR produces more empirical evidence, insurance companies will act accordingly. Indeed, occupational insurance for medical practitioners now specifically excludes any medical conditions that arise from exposure to non-ionizing radiation such as RF-EMF, including that from phones and mobile devices. Indeed, expect а strona response from the insurance industry as its actuaries evaluate the risks posed by long-term exposure to RFR in the weight of recent scientific evidence.

Regulators, such as the Securities and Exchange Commission, also recognise the economic impact of risks, as do mobile phone and internet services providers. Take, for example, that Vodafone and Verizon, among others, now include, and make provision for in their financial reports, the risks of litigation in relation to the health effects of products and services involving RFR, whether from smartphones or WiFi routers. Clearly neither believes that the small print in the safety information issued with RFR devices is sufficient. Take, for example, the following excerpt from Vodafone Group PLC, Annual Report.

"7. Our business may be impacted by actual or perceived health risks associated with the transmission of radio waves from mobile telephones, transmitters and associated equipment. Risk: Concerns have been expressed that the electromagnetic signals emitted by mobile telephone handsets and base stations may pose health risks. We are not aware that such health risks have been substantiated, however, in the event of a major scientific finding supporting this view this might result in prohibitive legislation being introduced by governments (or the European Union), a major reduction in mobile phone usage (especially by children), a requirement to move base station sites, significant difficulty renewing or acquiring site leases, and/or major litigation. An inadequate response to electromagnetic fields ('EMF') issues may result in loss of confidence in the industry and Vodafone."1

#### Discussion

In a submission to the United Nations in 2015, over 200 scientists requested it to address "the emerging public health crisis" related to the use of RFR emitting devices.46 They urged the United Nations Environmental Programme (UNEP) to review current exposure standards and to identify measures to substantially lower human exposures to microwave radiation. The scientists argued that existing "quidelines do not cover longterm exposure and low-intensity effects" and are "insufficient to protect public health." They note the urgency in this, as children are more vulnerable to the effects of RFR.

Microwave radiation is considered by majority of informed scientists as an invisible source of potentially toxic

<sup>&</sup>lt;sup>1</sup> Vodafone Group PLC, Annual Report 2012. https://www.vodafone.com/content /annualreport/annual\_report12/downloads/perform ance\_vodafone\_ar2012\_sections/principal\_risk\_fac tors\_and\_uncertainties\_vodafone\_ar2012.pdf.

pollution that scientific research in the sciences has identified as being harmful to biological systems and, ultimately, human health and well-being. Think of a smoke-filled bar of yore, where smokers and non-smokers alike are subjected to toxic carcinogens. Now, think of that same bar in countries where smoking is banned from such premises. However, have we replaced one hazard with another, if one considers the RFR being emitted by the WiFi routers, and radio units all of the smart devices in pubs, cafes, restaurants, homes, schools, and the workplace in the age of the Internet of Things (IoT), the scale of the dilemma that we have unthinkingly drifted into becomes clear. That, is of course, if one accepts the scientific evidence.

All this is of concern to informed computer scientists and technologists, who find the exposure to a multiplicity of, and close proximity to, WiFi signals problematic. Take, for example, Ajay Malik, SVP of Engineering and Products, Network World, who has also called for the WiFi standard to be reviewed by the FCC. He argues that the "amount of radiation exposure today is over 100 times higher as we live in proximity to a very large number of actively transmitting Wi-Fi Devices and Wi-Fi Access Points/Routers." 47 He therefore raises questions on the cumulative impact on adults and children of these unplanned for levels of exposure that often can go beyond SAR safety limits. Of course, he is unaware of the nonthermal health effects which perhaps, of greater concern, as the relevant mechanisms operate at lower exposure levels and durations.


The child in the image below is not operating his tablet device safely—that is, he is not in compliance with existing safety standards, inappropriate as they may be. As a consequence, his vital organs, eyes and brain are being exposed to unacceptable and potentially unsafe levels of microwave radiation. Over time the cells in his body will develop oxidative stress, due to elevated

levels of ROS and attenuated levels of anti-oxidants associated with exposure to microwave radiation. However, the bright light shining on his face is also affecting his developing eyes, which are more sensitive to those of an adult.

Vision issues aside, this light acts to significantly attenuate melatonin production in the brain. The first order effect here is interference with the circadian rhythm and sleep disturbance. The research literature on the effect that screens have in suppressing melatonin is levels unequivocal. Cajochen et al. provide convincing evidence of the effect that "A 5-h evening exposure to a white LED backlit screen... elicited a significant suppression of the evening rise in endogenous melatonin and subjective as well as objective sleepiness, as indexed by a reduced incidence of slow movements and EEG low-frequency activity (1-7 Hz) in frontal brain reaions." 48 Sleep disruption is also problematic as "sleep mediates learning and memory processing" and is vital for memory "encoding, consolidation, and reconsolidation, into the constellation of additional processes that are critical for development." 49 efficient memory However, as melatonin is also one of the body's most effective antioxidants and ROS scavengers, it is putting him at particular risk of second-order effects. It specifically increases the probability, low



What's wrong with this picture?



The mechanisms involve altered activity of antioxidant enzymes, altered immune system response, increased free radical accumulation in cells, oxidative damage to DNA, and apoptosis.

Figure 2 RFR Mechanisms and Outcomes

that this may be, that at some point in the future he may develop cancer as an adult. One must also consider the remote probability that he may develop cancers or other ill-effects conditions in childhood. However, scientific experiments have also demonstrated that exposure to WiFi radiation also affects brain development in young rats and their ability to learn and engage in

routine problem solving <sup>50</sup>, <sup>51</sup>, <sup>42</sup>. The implications for brain development in children is clear, as are the consequences for their immediate wellbeing.

#### **Conclusions**

As far back as 1973, a review and study by Russian scientists on the effects of low-intensity RFR on experimental animals indicated clear evidence of effects on the brain and nervous system, and also the heart and testes, of subjects. 52 Historically, Russia has more stringent safety standards than the West, whether it is the EU or US, when it comes to RFR.43 The evidence provided Russian scientists and contemporaries in the US and Europe should have given pause to telecommunications industry and regulators relation the in to commericalisation and widespread use of mobile telephony in the 1980s. However, in 2019 the cumulative body of scientific evidence should have governments and regulators take immediate action to change policy and implement appropriate safety standards for digital technologies, as it is children that are most at risk.

Given the clear risks that RFR-based technologies present, it is also vital for parents and educators to take immediate action on the use of microwave emitting devices where children are concerned. As there is overwhelming evidence that safety standards are woefully outdated, the action to be taken is clear. The precautionary principle should be applied and the use of all microwave RFRdevices, from WiFi-enabled enabled tablets (and smartphones) to WiFi routers, should be heavily curtailed or eliminated. The figure above summarizes paper's findings and provides compelling reasons for why such action is necessary.

As indicated, Figure 2 summarises the evidence of risk and indicates the role of specific mechanisms in producing the various threats to human health and well-being. Each of the outcomes identified are independent of each other; hence, the risk of some form of ill-health to children due to RFR exposure is highly probable.

At the risk of repetition, there is only one realistic course of action. Children and adolescents should not be using smartphones, or WiFi-enabled tablet

devices, and their expose to RFR sources should be minimized. This might seem impractical in the digital world, but in our real analogue world, children and teenagers are no longer permitted legal access to cigarettes, nor is it socially acceptable for adults to smoke in their presence. Given the current scientific evidence, the pathophysiological properties of RFR appear to be no different than cigarette smoke or similar carcinogens.

Thus, in light of the evidence, the precautionary principle should be applied governments should implement policies that result in the removal of WiFi routers and all WiFi devices from the classrooms of elementary/primary and secondary/high schools. Just to remind the reader what the precautionary *principle* means: "When an activity raises threats of harm to human health the environment, precautionary measures should be taken even if some cause and effect relationships are not fully established scientifically."2 We are well beyond that point, as this paper illustrates. The application of the precautionary principle is a statutory requirement in some areas of law in the European Union, as expressed in the Charter of Fundamental Rights. Thus EU governments at least have a political and an ethical responsibility to act.

absence of the appropriate government policy, educators need to reconsider the untrammeled use of WiFi in schools and not employ iPads or tablets for use by children in class. Devices that use e-Ink, or similar types of electronic paper display, as opposed to LED screens, should be used in the classroom and at home to access ebooks/texts, but these should operated in airplane mode when reading.

Parents and guardians also need to act and should consider the following

<sup>&</sup>lt;sup>2</sup> https://en.wikipedia.org/wiki/Precautionary\_principle

recommendations in order to exercise their personal duty of care:

- Educate children and adolescents about the health risks of all smart devices.
- Restrict device time to 30 minutes for all RFR-enabled devices, not just screen time.
- In respect of screen time, all LED screen devices should have a Blue Light Filter. Apps like F-Lux are ideal here. This minimises melatonin reduction in users.
- Smart phones have 2/3/4G, WiFi and Bluetooth radio units all of which are normally switched on. These should be used only when required. In addition, the small print on Health and Safety information that comes with a smartphone typically indicates that they should NOT be carried nor operated less than 2.5 cm from the body.
- Remember that the WiFi Safety standards for ALL devices is that they must be operated 20 cm or 8" from the body and for no more than 30 minutes.
  - Given the risk handing a young child an active RFR device, such as а smartphone or an iPad, to hold their in car is seat/pram, for all intents and purposes the same as giving them a cigarette to smoke.
- If children or adolescents have access to smartphones and WiFi devices, the devices should not be carried or operated on or near their person.
  - Ear & microphones should be used for all calls.

- If children are using screen device for games, they should be operated in airplane mode.
- Ensure WiFi routers are not in or near or directly beneath children's bedrooms and they should be switched off at night. No RFR device, including some types of baby monitors, should in in a child's bedroom.
- Minimise the use in the home of all Internet of Things (IoT) devices such as Smart Meters, Virtual Assistants, Hive, Chromecast, WiFi dongles, and so on.

There is also a clear onus on scientists and practitioners in the computing and IT industry to act and ensure that the safety standards for all RFR and WiFi devices are reviewed in light of the scientific findings. To otherwise would be irresponsible. There will be enormous resistance to change from vested interests and the political establishment. This has already happened, with orchestrated campaigns against natural scientists conducting independent research on the health implications of RFR, particularly in the US. An excerpt from a recent article in The Guardian newspaper summarises the type of response to be expected from industry with respect to microwave RFR and in particular the release of the findings of the NTP study. "Central to keeping the scientific argument going is making it appear that not all scientists agree. Towards that end, and again like the tobacco and fossil-fuel industries, the wireless industry has "war-gamed" science, as a Motorola internal memo in 1994 phrased it. War-gaming science involves playing offence as well as defence - funding studies friendly to the industry while attacking studies that raise questions; placing industry-friendly experts on advisory bodies such as the World Health Organisation and seeking

to discredit scientists whose views differ from the industry's."53

Returning to the quote at the beginning of this paper by Professor Frentzel-Beyme MD, we have, as the evidence adduced herein indicates, far exceeded the "level of proof required to justify action for health protection." The theory that non-ionizing RFR exposure could **not** cause cancer has been refuted using the scientific method. It is ironic, in the era of neoconservatism, neoliberalism, and the anti-environmental policies of cheerleader-in-chief Donald necon Trump, that the smoking gun should be provided by the National Toxicology Program of the US Department of Health and Human Services. This study, as indicated above, is just the latest of many to provide the "clear evidence" required for policy and social change.

The need for social change is this area is as important, and no less controversial, than that required to respond to the challenge of global warming. However, the forces resisting change to the status quo are considerable. Take for example that "Not one major news organisation in the US or Europe reported [the] scientific news [published by the NTP]. But then, news coverage of mobile phone safety has long reflected the outlook of the wireless industry."53 In order to combat vested interests and protect children, parents and grandparents, aunts and uncles, need to act to change extant social perspectives on seeminaly harmless digital technologies that entertain and beguile, and which offer without affordances apparent consequences. That will be the challenge for readers of this paper. To understand that technology is not neutral—that it has negative as well as positive consequences for users and society, and that there is a dark side to the bright screen on which you may be reading this article.

#### **About the Author**

Professor Tom Butler is a social scientist at University College Cork. A former satellite and microwave communications engineer and IT professional, he is more than familiar with the traditional safety issues relating to microwave RFR. His Pauline conversion from the engineering perspective on RFR thermal safety occurred through research engagements with the Chief Risk Officer of a global corporation who pointed out significant risks to children from the nonthermal effects of RFR. These discussions and related events in his personal life stimulated Professor Butler's interest in this important topic.

In a research context, Professor Butler is former Government of Ireland Research Fellow, Principal Investigator (PI) of the Governance Risk and Compliance Technology Centre (2013-2018), PI of the SmaRT and SamRT4Reg Commercialisation Fund Projects (2017-2019), and Co-PI of two Marie Skłodowska-Curie Career-FIT Fellowships in Artificial Intelligence (2019-2022). With over €8.5 million in research funding on the application of digital technologies to date, he has 220 publications and 11 inventions. Tom is a member of the European Commission's Expert Group on Regulatory Obstacles to Financial Innovation (ROFIEG) in the area of FinTech, a member of the Global RegTech Council, and a member of the Financial Industry Enterprise Data Management Council (EDMC).

#### References

<sup>&</sup>lt;sup>1</sup> National Toxicology Programme (2018). Cell Phone Radio Frequency Radiation Studies. https://www.niehs.nih.gov/health/materials/cell \_phone\_radiofrequency\_radiation\_studies\_508.

<sup>&</sup>lt;sup>2</sup> National Toxicology Programme (2018). High Exposure to Radio Frequency Radiation Associated With Cancer in Male Rats, Telephone Press Conference, 10/31/18, 2:00

- pm ET. https://www.nih.gov/news-events/news-releases/high-exposure-radio-frequency-radiation-associated-cancer-malerats.
- <sup>3</sup> Food and Drug Administration (FDA) (1999). Nomination Letter to Coordinator of NTP Chemical Nomination and Selection Committee. nomihttps://ntp.niehs.nih.gov/ ntp/htdocs/chem\_background/exsumpdf/wirele ss051999\_508.pdf.
- <sup>4</sup> Vijayalaxmi & Obe, G., (2004). Controversial cytogenetic observations in mammalian somatic cellsexposed to radiofrequency radiation, Radiat. Res. 162 (5), 481–96.
- Morgan, L. L., Miller, A. B., Sasco, A., & Davis, D. L. (2015). Mobile phone radiation causes brain tumors and should be classified as a probable human carcinogen (2A). International journal of oncology, 46(5), 1865-1871.
- 6 http://ehtrust.org/the-california-medicalassociation-wireless-resolution/
- <sup>7</sup> https://ecfsapi.fcc.gov/file/7520941318.pdf
- <sup>8</sup> https://www.nytimes.com/2018/11/01/health/cellphone-radiation-cancer.html
- <sup>9</sup> Melnick, R. L. (2019). Commentary on the utility of the National Toxicology Program Study on cell phone radiofrequency radiation data for assessing human health risks despite unfounded criticisms aimed at minimizing the findings of adverse health effects. Environmental research, 168, 1-6.
- <sup>10</sup> https://www.ramazzini.org/comunicato/ondeelettromagnetiche-listituto-ramazzini-rispondeallicnirp/
- Falcioni, L., Bua, L., Tibaldi, E., Lauriola, M., De Angelis, L., Gnudi, F., Mandrioli, D., Manservigi, M., Manservisi, F., Manzoli, I. & Menghetti, I. (2018). Report of final results regarding brain and heart tumors in Sprague-Dawley rats exposed from prenatal life until natural death to mobile phone radiofrequency field representative of a 1.8 GHz GSM base station environmental emission. Environmental research, 165, 496-503.
- Wyde, M. (2016). NTP toxicology and carcinogenicity studies of cell phone radiofrequency radiation. BioEM2016 Meeting, Ghent, Belgium. https://ntp.niehs.nih.gov/ntp/research/areas/cell phone/slides\_bioem\_wyde.pdf.
- <sup>13</sup> Philips, A.. Henshaw, D., L Lamburn, G. & M. O'Carroll, (2018). Brain tumours: rise in Glioblastoma Multiforme incidence in England 1995–2015 suggests an adverse environmental

- or lifestyle factor, Journal of Environmental and Public Health, vol. 2018, Article ID 7910754.
- <sup>14</sup> Röösli, M., Lagorio, S., Schoemaker, M. J., Schüz, J., & Feychting, M. (2019). Brain and Salivary Gland Tumors and Mobile Phone Use: Evaluating the Evidence from Various Epidemiological Study Designs. Annual review of public health, 40.
- <sup>15</sup> Coureau, G., Bouvier, G., Lebailly, P., Fabbro-Peray, P., Gruber, A., Leffondre, K., ... & Baldi, I. (2014). Mobile phone use and brain tumours in the CERENAT case-control study. Occup Environ Med, oemed-2013.
- <sup>16</sup> Mialon, H. M., & Nesson, E. T. (2018). Mobile Phones and the Risk of Brain Cancer Mortality: A Twenty-Five Year Cross-Country Analysis.
- <sup>17</sup>Kabbat, G.(2017). Are brain cancer rates increasing? https://www.forbes.com/sites/ geoffreykabat/ 2017/12/23/are-brain-cancerrates-increasing-and-do-changes-relate-to-cellphone-use/#12df118e25b4
- West, J. G., Kapoor, N. S., Liao, S. Y., Chen, J. W., Bailey, L., & Nagourney, R. A. (2013). Multifocal breast cancer in young women with prolonged contact between their breasts and their cellular phones. Case reports in medicine, 2013.
- <sup>19</sup> Chauhan, P., Verma, H. N., Sisodia, R., & Kesari, K. K. (2017). Microwave radiation (2.45 GHz)-induced oxidative stress: Wholebody exposure effect on histopathology of Wistar rats. Electromagnetic Biology and Medicine, 36(1), 20-30.
- <sup>20</sup> Carpenter, D. O., & Bushkin-Bedient, S. (2013). Exposure to chemicals and radiation during childhood and risk for cancer later in life. Journal of Adolescent Health, 52(5), S21-S29.
- <sup>21</sup> Christ, Andreas, Marie-Christine Gosselin, Maria Christopoulou, Sven Kühn, & Niels Kuster. (2010). Age-dependent tissue-specific exposure of cell phone users. *Physics in medicine and biology*, 55(7): 1767.
- <sup>22</sup> Keshvari, J., Keshvari, R., & Lang, S. (2006). The effect of increase in dielectric values on specific absorption rate (SAR) in eye and head tissues following 900, 1800 and 2450 MHz radio frequency (RF) exposure. *Physics in Medicine and Biology*, 51(6), 1463.
- <sup>23</sup> Gandhi, O. P., Morgan, L. L., de Salles, A. A., Han, Y. Y., Herberman, R. B., & Davis, D. L. (2012). Exposure limits: the underestimation of absorbed cell phone radiation, especially in

- children. Electromagnetic Biology and Medicine, 31(1), 34-51
- <sup>24</sup> Portier, C.J., & Leonard W.L. (2016). Do Cell Phones Cause Cancer? Probably, but It's Complicated, Scientific American, June 13, 2016.
- <sup>25</sup> Frentzel-Beyme, R. (1994). John R. Goldsmith on the usefulness of epidemiological data to identify links between point sources of radiation and disease. Public health reviews, 22(3-4), 305-320.
- Waris G, Ahsan H. (2016). Reactive oxygen species: role in the development of cancer and various chronic conditions. Journal of Carcinogenesis. 2006;5(14). doi:10.1186/1477-3163-5-14.
- <sup>27</sup> De Iuliis, G. N., Newey, R. J., King, B. V., & Aitken, R. J. (2009). Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. PloS one, 4(7), e6446.
- Yakymenko, I., Tsybulin, O., Sidorik, E., Henshel, D., Kyrylenko, O., & Kyrylenko, S. (2016). Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. Electromagnetic biology and medicine, 35(2), 186-202.
- <sup>29</sup> Nazıroğlu, M., Yüksel, M., Köse, S. A., & Özkaya, M. O. (2013). Recent reports of Wi-Fi and mobile phone-induced radiation on oxidative stress and reproductive signaling pathways in females and males. The Journal of membrane biology, 246(12), 869-875.
- <sup>30</sup> Pall, M. L. (2018). Wi-Fi is an important threat to human health. Environmental research, 164, 405-416.
- <sup>31</sup> Pall, M.L., (2013). Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. J. Cell. Mol. Med. (17), 958–965. http://dx.doi.org/10.1111/jcmm.12088.
- <sup>32</sup> Kesari, K. K., Siddiqui, M. H., Meena, R., Verma, H. N., & Kumar, S. (2013). Cell phone radiation exposure on brain and associated biological systems. Indian Journal of Experimental Biology, March 2013; 51: 187– 200.
- <sup>33</sup> Han, Y. Y., Gandhi, O. P., De Salles, A., Herberman, R. B., & Davis, D. L. (2010). Comparative assessment of models of electromagnetic absorption of the head for children and adults indicates the need for policy changes. Eur J Oncol. Volume, 5, 301-318.

- <sup>34</sup> Kheifets, L., Repacholi, M., Saunders, R., & Van Deventer, E. (2005). The sensitivity of children to electromagnetic fields. Pediatrics, 116(2), e303-e313.
- <sup>35</sup> Akhavan-Sigari, R., Baf, M. M. F., Ariabod, V., Rohde, V., & Rahighi, S. (2014). Connection between cell phone use, p53 gene expression in different zones of glioblastoma multiforme and survival prognoses. Rare Tumors, 6(3).
- <sup>36</sup> Hardell, L., & Carlberg, M. (2015a). Increasing rates of brain tumours in the Swedish national inpatient register and the causes of death register. International journal of environmental research and public health, 12(4), 3793-3813.
- <sup>37</sup> Hardell, L., & Carlberg, M. (2015b). Mobile phone and cordless phone use and the risk for glioma–Analysis of pooled case-control studies in Sweden, 1997–2003 and 2007–2009. Pathophysiology, 22(1), 1-13.
- <sup>38</sup> Carlberg, M., Hedendahl, L., Ahonen, M., Koppel, T., & Hardell, L. (2016). Increasing incidence of thyroid cancer in the Nordic countries with main focus on Swedish data. BMC cancer, 16(1), 426.
- <sup>39</sup> Tillmann, T., Ernst, H., Streckert, J., Zhou, Y., Taugner, F., Hansen, V., Dasenbrock, C., (2010). Indication of cocarcinogenic potential of chronic UMTS-modulated radiofrequency exposure in an ethylnitrosourea mouse model. Int. J. Radiat. Biol. 86, 529–541.
- <sup>40</sup> Lerchl, A., Klose, M., Grote, K., Wilhelm, A. F., Spathmann, O., Fiedler, T., ... & Clemens, M. (2015). Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. Biochemical and biophysical research communications, 459(4), 585-590.
- <sup>41</sup>https://drive.google.com/file/d/0B14R6QNkmaXu d2dQZmtJZWpKOW8/view. Accessed February 2019.
- <sup>42</sup> Belpomme, D., Hardell, L., Belyaev, I., Burgio, E., & Carpenter, D. O. (2018). Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective. Environmental pollution, 242, 643-658.
- <sup>43</sup> Grigoriev Y. (2017). Methodology of Standards Development for EMF RF in Russia and by International Commissions: Distinctions in Approaches. In Markov, M (Ed.), Dosimetry in Bioelectromagnetics. Chapter 15. pp. 315-337. Boca Raton, FL: Taylor & Francis.
- <sup>44</sup> Hankin, N. (2002). Letter from Norbert Hankin, U.S. Environmental Protection Agency, to

Janet Newton President, The EMR Network (July 16, 2002), available at http://www.emrpolicy.org/litigation/case\_law/docs/noi\_epa\_response.pdf .Accessed 21-03-2017.

<sup>45</sup>https://www.lloyds.com/~/media/lloyds/reports/e merging%20risk%20reports/emf%20final%20 november%202010.pdf.

46 https://emfscientist.org/

- <sup>47</sup> http://www.networkworld.com/article/3004112/ wi-fi/why-the-fccs-safety-guidelines-for-wi-fineed-to-be-re-evaluated.html
- <sup>48</sup> Cajochen, C., Frey, S., Anders, D., Späti, J., Bues, M., Pross, A., Mager, R., WirzJustice, A., & Stefani, O., (2011). Evening exposure to a light-emitting diodes (LED)- backlit computer screen affects circadian physiology and cognitive performance. Journal of Applied Physiology 110, 1432e1438.
- <sup>49</sup> Walker, M. P., and Stickgold, R. (2004). Sleep-dependent learning and memory consolidation. Neuron 44, 121–133. doi: 10.1016/j.neuron.2004.08.031
- <sup>50</sup> Ikinci, A., Odaci, E., Yildirim, M., Kaya, H., Akça, M., Hanci, H., ... & Bas, O. (2013). The effects of prenatal exposure to a 900 megahertz electromagnetic field on hippocampus morphology and learning behavior in rat pups. NeuroQuantology, 11(4), 582-590.
- Narayanan, S. N., Kumar, R. S., Karun, K. M., Nayak, S. B., & Bhat, P. G. (2015). Possible cause for altered spatial cognition of prepubescent rats exposed to chronic radiofrequency electromagnetic radiation. Metabolic brain disease, 30(5), 1193-1206.
- <sup>52</sup> Tolgskaya, M.S. & Gordon, Z.V., (1973). Pathological Effects of Radio Waves, Translated from Russian by B Haigh. Consultants Bureau, New York/London 1-146.
- <sup>53</sup> Hertsgaard, M & Dowie, M. (2018). The inconvenient truth about cancer and mobile phones. The Guardian, Sat 14 Jul 2018 15.00 BST. <a href="https://www.theguardian.com/technology/">https://www.theguardian.com/technology/</a> 2018/jul/14/mobile-phones-cancer-inconvenient-truths. Accessed Sunday 17th February, 2019.